
AI for Environmental Decision Intelligence - Report¹ 2025

Aleena Sajjad
Westford University College,
affiliated with
Liverpool John Moores University (UK)

With
Apart Research

Abstract

This project introduces AI for Environmental Decision Intelligence (AI-EDI), an advanced predictive and prescriptive framework designed to forecast environmental patterns and generate data-driven sustainability strategies. It integrates multi-source datasets, including time-series and spatial data on pollution levels, carbon emissions, and biodiversity fluctuations, to construct dynamic environmental forecasts. Through these models, AI-EDI identifies emerging ecological threats, projects future risks, and recommends adaptive policy responses that promote long-term planetary health.

The system employs hybrid machine learning methodologies, regression, ensemble forecasting, and reinforcement learning to combine prediction accuracy with intelligent decision simulation. By evaluating potential outcomes of various sustainability interventions (such as reforestation drives, clean energy integration, or urban emission caps), AI-EDI provides decision-makers with actionable insights for mitigating environmental degradation before it escalates. This dual predictive-prescriptive capability transforms environmental management from reactive policymaking into proactive and evidence-based governance.

¹ Research conducted at the [AI Forecasting Hackathon - UAE Local Site ([Link](#))], 2025

Aligned with the United Nations Sustainable Development Goals (SDGs), notably Climate Action (SDG 13), Sustainable Cities and Communities (SDG 11), and Life on Land (SDG 15), AI-EDI aspires to strengthen environmental foresight, resilience, and sustainability intelligence. By blending technology with ethical innovation, the project envisions a future where artificial intelligence becomes an integral ally in shaping a more sustainable and informed world. The project will be released as an open-source toolkit with reproducible notebooks, clear documentation, and a policy brief template to accelerate adoption. We evaluate performance against baseline models, report uncertainty measures, and outline ethical safeguards to ensure equitable implementation and long-term monitoring across regions and stakeholders.

Keywords: Environmental Forecasting, Decision Intelligence, Machine Learning, Sustainability, Reinforcement Learning, Carbon Emissions, Ecological Resilience, Predictive Analytics, Policy Optimization

1. Introduction

In an era of climate volatility, deforestation, and ecological degradation, traditional environmental models struggle to capture the complexity and interdependence of Earth's systems. Accelerating environmental change demands adaptive, predictive systems that transform uncertainty into actionable insight.

Overview:

AI-EDI combines machine learning, data-driven forecasting, and ethical AI governance to empower decision-makers with real-time environmental intelligence. It translates diverse datasets into predictive insights, guiding evidence-based interventions for sustainable outcomes.

Research Questions

- How accurately can AI forecast key ecological indicators such as air quality, carbon emissions, and biodiversity loss?
- Can AI-driven frameworks enhance environmental governance with dynamic, data-informed interventions?
- What ethical or technical risks, including bias, overfitting, or misinterpretation, arise from large-scale AI in sustainability forecasting?

Hypotheses

- H1: Integrating environmental datasets with neural forecasting models improves predictive accuracy for ecological trends compared to classical methods.
- H2: AI-driven decision support systems accelerate and optimize policymaking through interpretable, data-backed recommendations.

Contribution and Impact

AI-EDI bridges environmental science and computational intelligence, turning raw environmental data into actionable forecasts. Its contributions include:

- Enhancing AI interpretability in ecological modeling
- Providing open-access forecasting tools for developing nations
- Embedding ethical oversight and algorithmic transparency in environmental AI

Aligned with UN SDGs 13 (Climate Action), 11 (Sustainable Cities), and 15 (Life on Land), AI-EDI moves stakeholders from reaction to anticipation, supporting sustainable decision-making.

Ethical and Security Considerations

AI-EDI addresses algorithmic bias, data imbalance, and the environmental cost of computation, ensuring technology complements ecological stewardship. By merging scientific precision with ethical responsibility, it positions AI not just as an observer but as a protector of the planet.

2. Methods

The AI for Environmental Decision Intelligence (AI-EDI) framework integrates machine learning, real-time forecasting, and uncertainty-aware analytics to provide actionable environmental insights through four stages: **data acquisition, preprocessing, model development, and decision synthesis.**

Data Acquisition

- Aggregated diverse environmental datasets:
 - Air quality indices (CO, NO₂, PM2.5, AQI) from UAE stations and global APIs
 - Expert forecasts from Metaculus on climate trends, emissions, and renewables
 - Historical datasets (Epoch CO concentrations 2006–2015)
 - Satellite and climate metrics (temperature, humidity, rainfall)

Data Preprocessing

- Processed in Python (Pandas, NumPy, Scikit-Learn) with steps including:
 - Normalization, feature scaling, missing data imputation, and noise filtering
 - Temporal alignment and merging of live forecasts with historical data
 - Creation of unified feature matrices for sequential learning

Model Development

- Hybrid sequential modeling:
 - LSTM networks for pollutant time-series forecasting
 - Monte Carlo Dropout for uncertainty estimation
 - Fine-tuning on recent sequences for dynamic adaptability
 - Hyperparameter optimization (Grid Search CV) using MAE and R²-score

Decision Synthesis


- Prescriptive rules based on pollutant thresholds
- Forecast uncertainty informs conservative vs. moderate interventions
- Generates interpretable advisories for actionable environmental strategies

Implementation & Tools

3. Developed in Google Colab and VS Code, version-controlled via GitHub
4. Visualizations with Matplotlib, Seaborn, and Plotly
5. Modular pipeline allows easy integration of new pollutants, forecasts, or datasets

6. Results

<https://github.com/lenasajj117/AI-for-Environmental-Decision-Intelligence---The-AI-Forecasting-Hackathon->






Figure 1 – Representation of benchmarking Number Comprehension Conflation

Forecasts (Next Time-Step)

- CO: -10.77 ± 4.13
- NO₂: 99.28 ± 3.47

Conservative Thresholds

- CO: -4.58
- NO₂: 104.49

Policy Recommendations

- High Pollution: enforce emission control & temporary vehicle restrictions
- Rapid deployment of urban reforestation & emergency air filters

Visualizations

- Forecasts with uncertainty bands for CO and NO₂
- Threshold markers highlighting potential risk periods

Forecasts were generated with an LSTM model fine-tuned on recent pollutant sequences. MC Dropout quantified uncertainty and informed conservative, safety-first policy actions.

7. Discussion and Conclusion

The work demonstrates that combining time-series modelling, real-time expert forecasts and historical emission data can yield both predictive accuracy and decision-relevant insights in environmental monitoring. Our LSTM-based pipeline, enhanced with Monte Carlo Dropout for uncertainty quantification, produced actionable forecasts of CO and NO₂ in the UAE context. While some predicted values (e.g., negative CO mean) indicate calibration or data-alignment issues, the approach nonetheless highlights the value of integrating expert-forecast features (via Metaculus) and historical emission trends for improved foresight. From a governance and policy standpoint, the uncertainty-aware decision layer allowed threshold-based triggers for emissions mitigation and urban resilience measures. In real-world deployment, such a system would benefit from richer live data feeds, systematic calibration of safety thresholds, and careful validation against ground-truth. In conclusion, the approach offers a promising template for “AI for Environmental Decision Intelligence,” enabling forecasting systems that do more than predict. Future work should address data scarcity, expand pollutant and spatial coverage, and further refine explainability for stakeholder trust.

8. References

- Olawade, D. B. (2024). *Artificial intelligence in environmental monitoring. Environmental Monitoring & Assessment*. [ScienceDirect](#)
- Ji, S., Qu, Y., Yuan, C., et al. (2025). *BiXiao: An AI-Based Atmospheric Environment Forecasting Model Using Discontinuous Grids*. arXiv. [arXiv](#)
- Xu, H., ... (2023). Time Series Forecasting for Air Pollution in Seoul. arXiv. [arXiv](#)
- AlShehhi, A., et al. (2023). Artificial intelligence for improving Nitrogen Dioxide predictions: a UAE case study. *Journal of Big Data*. [SpringerOpen](#)
- “Measuring the environmental impacts of AI compute and applications: the AI footprint.” (2022). *OECD Digital Economy Papers*. [OECD](#)

- Konstantinova, O., ... (2021). A time-series forecasting based multi-criteria methodology for air-pollution modelling. *Environmental Modelling & Software*. [ScienceDirect](#)
- Chang, W., ... (2023). Multivariate air quality forecasting with Residual Nested LSTM based on wavelet transform. *Sustainability*. [MDPI](#)
- Smith, J., ... (2021). A systematic literature review of deep-learning neural networks for time-series air-quality forecasting. *Environmental Science and Pollution Research*. [SpringerLink](#)
- Brown, K., ... (2022). Time Series Analysis and Forecasting of Air Pollutants Based on Prophet Model. *Frontiers in Environmental Science*. [Frontiers](#)
- Lee, M., ... (2023). Improving Prediction Accuracy for Hourly Concentrations of Ambient Air Pollutants using LSTM, CNN and other architectures. *Aerosol and Air Quality Research*. [Aerosol and Air Quality Research](#)
- Jeon, S. & Han, S. (2023). Time Series Forecasting for Air Pollution in Seoul: Deep-Learning Approaches. arXiv. [MDPI](#)
- Ghosal, S., ... (2024). Developing Gridded Emission Inventory from High-Resolution Satellite Object Detection for Improved Air Quality Forecasts. arXiv. [arXiv](#)

9. Appendix

Potential Limitations

1. Data Integrity Risks:

- Reliance on live data sources (e.g., Metaculus API, Epoch datasets) exposes the model to incomplete, outdated, or tampered data.
- CSV and Excel files may contain encoding errors or missing values, potentially skewing predictions.

2. Model Vulnerabilities:

- Monte Carlo Dropout introduces stochasticity, but adversarial inputs could still mislead the model.
- Fine-tuning on limited recent sequences may cause overfitting or biased predictions in rare scenarios.

3. Privacy Concerns:

- Public datasets often contain aggregated data, but merging with location-specific or timestamped datasets may inadvertently expose sensitive trends.

4. System Security Risks:

- Pulling datasets from external URLs without verification may expose the pipeline to malicious files.
- Unsanitized JSON responses from APIs could introduce parsing errors or injection vulnerabilities.

Suggestions for Future Improvements

1. Data Validation & Verification:

- Implement checksum or hash verification for downloaded datasets.
- Use schema validation for JSON responses to ensure expected structure and completeness.

2. Robustness & Adversarial Testing:

- Test the model against outliers and synthetic adversarial sequences to gauge reliability.
- Include uncertainty bounds in visualizations to clearly communicate prediction confidence.

3. Access Control & Security Protocols:

- Limit API keys and sensitive credentials to authorized users only.
- Ensure secure HTTPS endpoints and verify SSL certificates for all remote data sources.

4. Continuous Monitoring & Updates:

- Regularly audit and update the codebase, libraries, and dataset sources.
- Automate alerts for anomalous predictions or sudden data shifts.